

Mouth care after surgery



Information for patients

This leaflet aims to help you understand what you can expect after having mouth surgery and contains answers to many commonly asked questions. If you have any other questions or would like further explanation, please ask.

Confirming your identity

Before you have a treatment or procedure, our staff will ask you your **name** and **date of birth** and check your **ID band**. If you don't have an ID band we will also ask you to confirm your address.

If we don't ask these questions, then please ask us to check.

Ensuring your safety is our primary concern.

What did the surgery involve?

You will probably have had your gum cut and lifted away from the underlying bone. This is known as 'raising a flap'. It is done to uncover a root, or a tooth, which has been difficult to extract or to identify and to remove something in your jawbone, such as a small cyst. Some bone will probably have been drilled away and your gum will have been stitched up.

What should I do after the surgery?

- Do not rinse or spit for the rest of the day because this can make your wound bleed. Also, do not suck or put your tongue into the hole.
- The next day, gently rinse out your mouth using warm, salty water or a mouthwash. In particular, do this after you eat to prevent food sticking to your stitches or wound.
- Avoid physical exertion for the first 24 hours because this can also make your wound bleed.
- Take your painkillers as directed by your dentist.

What problems might occur?

- Your wound will be sore for two or three days. Please take the pain-killing drugs your dentist has prescribed for you as directed.
- Infection is rare. You will not normally be prescribed antibiotics.
- You may have a slight oozing from your wound and a bad taste. This often happens for a few days but does not mean there is a problem.
- If your wound bleeds, roll up some gauze (not tissue) or a clean handkerchief into a pad, place over the area that is bleeding and bite down firmly for 10 minutes. If you are still bleeding when you remove the pad, repeat this procedure for 15 minutes. If you are still bleeding after doing this, please contact the department and arrange to return so that we can help you. If this happens at night or at the weekend, please ring **NHS 111** or your own General Dental Practitioner.
- Your teeth next to where you have had your treatment might feel tender for a few days. This is normal.



- You might have some facial swelling over the operation site and after some procedures you might notice that the feeling in your skin alters slightly.
- If you have had an operation on your lower jaw it is likely to be stiff and you will find it difficult to open your mouth. This will gradually improve but might take up to three weeks.

Is there anything else I should know?

- You will be given dissolvable stitches which do not need to be removed. They might take up to two weeks to dissolve, sometimes longer.
- Your saliva and tongue will be speckled with blood, which is normal.
- Be careful when you are eating or drinking hot drinks while your face and gums are still numb so you do not accidentally bite or burn your lips.

Who can I contact with queries and concerns?

Monday to Friday between 9am and 4.30pm: Tel: **020 3299 3480**

Department of Community Special Care Dentistry

King's College London Dental Institute

King's College Hospital, Bessemer Road, London SE5 9RS

After 4.30pm and at weekends please ring **NHS 111** or your **General Dental Practitioner**.

Sharing your information

We have teamed up with Guy's and St Thomas' Hospitals in a partnership known as King's Health Partners Academic Health Sciences Centre. We are working together to give our patients the best possible care, so you might find we invite you for appointments at Guy's or St Thomas'. To make sure everyone you meet always has the most up-to-



date information about your health, we may share information about you between the hospitals.

Care provided by students

We provide clinical training where our students get practical experience by treating patients. Please tell your doctor or nurse if you do not want students to be involved in your care. Your treatment will not be affected by your decision.

PALS

The Patient Advice and Liaison Service (PALS) is a service that offers support, information and assistance to patients, relatives and visitors. They can also provide help and advice if you have a concern or complaint that staff have not been able to resolve for you. The PALS office is located on the ground floor of the Hambleton Wing, near the main entrance on Bessemer Road - staff will be happy to direct you.

Tel: **020 3299 3601**

Email: **kch-tr.PALS@nhs.net**

You can also contact us by using our online form at
www.kch.nhs.uk/contact/pals

If you would like the information in this leaflet in a different language or format, please contact PALS on 020 3299 1844.