

Caring for your child's mouth after dental treatment under general anaesthetic

Information for patients, parents and carers

This leaflet explains how to care for your child's mouth after dental treatment under general anaesthetic. If you have any questions, please speak to the dentist looking after your child.

On the day of the procedure

- Your child should rest for the remainder of the day. They may eat and drink as soon as they feel able, but soft and cold food is more comfortable (for example, yoghurt, smoothies, mashed banana, eggs and cool soup). Hot food should be avoided as it may cause bleeding to occur.
- Do not allow your child to rinse their mouth for 24 hours. This may dislodge the blood clot and cause bleeding.
- If bleeding does occur, put pressure on the socket by placing a tightly rolled-up damp cotton gauze or clean cotton handkerchief directly onto the socket. Ask your child to bite hard for at least 10 minutes without disturbing the clot. If this fails to control the bleeding, call the hospital (our contact details are below).
- If your child has been given local anaesthetic to numb the area, be careful that they do not chew the inside of their cheek or bite their lip while it remains numb.
- Avoid touching the wounds with fingers. Stitches are usually dissolvable, so you do not need to return to have them removed unless your surgeon has requested for you to do so.
- The teeth should be brushed as usual, at bedtime, on the day of the operation, but take care to avoid any extraction sites and not to dislodge any blood clots. Your child should not rinse their mouth after brushing.

The next few days

Warm salt mouth washes can be used the day following surgery. To make these, mix one level teaspoon of salt in a glass of warm water. Repeat three times daily for up to five days.

When can my child go back to school and exercise?

It is important that your child stays off school for at least one day after the operation. This is because your child's judgement, performance and reaction time may be affected by the anaesthetic.

We also recommend that your child does not do any vigorous exercise for at least 24 hours after their operation. They should remain in the immediate care of a responsible adult for 24 hours after their operation.

What should I do if my child has a problem?

If you have any further questions or concerns, or if your child has persistent bleeding, severe swelling or severe pain, please contact the hospital on the telephone numbers listed below. Out of hours and at weekends you should call NHS 111 or attend your local emergency department (A&E) for advice.

A rare side effect of general anaesthetic is urine retention (that is, when your child cannot pass urine). If your child has not or cannot pass urine, please bring them to your local A&E department.

Pain relief

If there is pain after the extractions, use whatever you normally give your child for pain relief. A suitable painkiller is sugar-free paracetamol or ibuprofen for children. Read the instructions on the packet carefully and be sure that the stated dose is not exceeded. Next medicines due:

Paracetamol	Ibuprofen

The contents of this leaflet have been adapted for use with permission from the Department of Paediatric Dentistry in Guy's and St Thomas' NHS Foundation Trust.

Contact us

If you have any questions or concerns about the information in this leaflet, please contact the Department of Paediatric Dentistry team. If your child has persistent bleeding, severe swelling or severe pain, please contact us. If out of hours or at weekends, please call NHS 111 or attend your local A&E department for advice.

Central Referral Office	020 3299 4988
Clinic Appointments	020 3299 3055
Secretary	020 3299 3375
	020 3299 4983

For more information leaflets on conditions, procedures, treatments and services offered at our hospitals, please visit www.kch.nhs.uk/patientsvisitors/patients/leaflets

Sharing your information

We have teamed up with Guy's and St Thomas' Hospitals in a partnership known as King's Health Partners Academic Health Sciences Centre. We are working together to give our patients

the best possible care, so you might find we invite you for appointments at Guy's or St Thomas'. To make sure everyone you meet always has the most up-to-date information about your health, we may share information about you between the hospitals.

Care provided by students

We provide clinical training where our students get practical experience by treating patients. Please tell your dentist if you do not want students to be involved in your care. Your treatment will not be affected by your decision.

PALS

The Patient Advice and Liaison Service (PALS) is a service that offers support, information and assistance to patients, relatives and visitors. They can also provide help and advice if you have a concern or complaint that staff have not been able to resolve for you. They can also pass on praise or thanks to our teams. The PALS office is located on the ground floor of the Hambleton Wing, near the main entrance on Bessemer Road - staff will be happy to direct you.

PALS at King's College Hospital, Denmark Hill, London SE5 9RS

Tel: 020 3299 3601

Email: kch-tr.palsdh@nhs.net

If you would like the information in this leaflet in a different language or format, please contact our Communications and Interpreting telephone line on 020 3299 4826 or email kch-tr.accessibility@nhs.net